Care of Your Pet Following Surgery or Hospitalization

Pets, like people, are made up of systems of delicate tissues. In order to minimize post-surgical complications, you will need to provide the proper conditions if we are to expect satisfactory healing of your pet.

	Exercise and Activity:
0	Your pet should be under strict confinement to a cage or small room for days. Carry outside for
	elimination. No running, jumping or access to stairs should be permitted.
0	Brief leash walks (on a short leash) only fordays. Avoid stairs and jumping on and off of
	furniture. Outdoor exercise must be limited to short periods on a short leash and only when necessary.
0	No bathing or grooming or swimming fordays.
0	It is okay to resume normal activity after
	Incision/Wound Care:
0	Check the incision twice daily and notify us if you notice any swelling, redness or drainage. Prevent licking, chewing, and / or scratching at the incision / bandage.
0	Use shredded news paper instead of clay littler for seven days for declaws.
0	Keep the bandage/splint clean and dry. Cover loosely with a plastic bag when outdoors. Call the
	hospital if the bandage/splint slips, becomes wet, soiled or if you notice any swelling above or below
	the bandage/splint.
Diet:	
0	Return to your pet's normal diet.
0	Feed softened or canned food for the next three (3) days to allow your pet's mouth to heal.
0	A permanent diet change to is necessary. In order for this diet to be effective, it is
	imperative that you feed only this food. No table scraps or non-prescription treats allowed.
	Medications:
0	No medications are indicated.
0	Give the prescribed medications as indicated. (directions attached):
	Appointments:
0	We need to see your pet back for a recheck exam indays. Please call ahead for an appointment.
0	Suture removal indays:
0	Drain removal indays:
0	Drain removal indays: Bandage/splint change/check indays:
Monitor:	
0	A decrease in activity or appetite for one or two days may be observed and patients may not have a

PLEASE CALL US AT (843) 744-8927 IF YOU HAVE ANY QUESTIONS OR CONCERNS

drink water over one day, (3) weakness, (4) depression, (5) vomiting, or (6) diarrhea.

bowel movement for 24 to 36hours after surgery. This is normal. However, if your pet exhibits any of the following symptoms, please notify the hospital: (1) loss of appetite for over two days, (2) refusal to